



A Bova
combo
formulation

GingiPro™

For Feline Chronic Gingivostomatitis (FCGS)

Doxycycline (as monohydrate)/EIDD-1931/Meloxicam
60 mg/204 mg/0.15 mg/mL
Chicken-flavored oral paste, 30 mL pump

Also available: **GingiPro Duo™** (Doxycycline as monohydrate/EIDD-1931) and
GingiPro Solo™ (EIDD-1931 alone), 30 mL pump



✓ Oral Paste

✓ 3-in-1 Drug

✓ Less Stress

Understanding FCGS

Feline Chronic Gingivostomatitis (FCGS) is a painful, immune-mediated inflammatory condition of the oral mucosa in cats. Unlike regular gingivitis, FCGS involves ulcerative or proliferative lesions extending beyond the gingiva to the caudal oral cavity and pharynx.

The Culprit: Feline Calicivirus (FCV): Research shows that FCGS is closely linked to FCV - a stubborn virus that causes persistent infection in the mouth. However, not all cats exposed to FCV develop this condition.

EMERGING THERAPY

For the treatment of FCV-associated FCGS

- 1 EIDD-1931**
Active antiviral metabolite targeting FCV RNA replication
- 2 Doxycycline (as monohydrate)**
Broad-spectrum antibiotic with immunomodulatory effects
- 3 Meloxicam**
NSAID to reduce inflammation and discomfort
- 4 Sustained improvement seen in most cats within 2-3 months**
- 5 Effective even in post-extraction cases and juvenile gingivitis**

Minimize invasive extractions.
This product can be used as part of the treatment of FCGS with precision—using a simple, targeted, and feline-appropriate solution**.



Scan to read: *Pathology of Feline Chronic Gingivostomatitis* by Dr. Richard Malik & Dr. Andrea Harvey

Prescribe for a patient today!
Online via iFill.com or call 800-754-5222

GingiPro™

Doxycycline (as monohydrate)/EIDD-1931/Meloxicam
60 mg/204 mg/0.15 mg/mL
Chicken flavored oral paste, 30 mL pump

Standard Protocol (Bova Dose Rate)

Doxycycline: 5 mg/kg BID
EIDD-1931: 17 mg/kg BID
Meloxicam: 0.0125 mg/kg BID

Administration

Best: Apply directly to oral mucosa for high local drug concentration. Full safety instructions are provided with the product.

Alternate: Can be mixed into wet food if direct dosing is not tolerated.

DOSING GUIDE (using Topi-CLICK Perl® dispenser)

Weight	3 kg cat	4.5 kg cat	6 kg cat
Total clicks per day	2 clicks	3 clicks	4 clicks
Frequency	Twice Daily	Twice Daily	Twice Daily
Daily Dosing	1 click in AM & 1 click in PM	2 clicks in AM 1 click in PM or 1 click in AM 2 clicks in PM	2 clicks in AM & 2 clicks in PM
Doxycycline/EIDD-1931/ Meloxicam (Strength per dose)	15 mg 51 mg 0.0375 mg	22.5 mg 76.5 mg 0.05625 mg	30 mg 102 mg 0.075 mg

Topi-CLICK Perl®
1 click = 0.25 mL

- EIDD-1931 is designed for immediate local action as an antiviral, unlike molnupiravir, which requires liver activation to become effective.
- The combination of three medications improves healing and immune regulation compared to using a single drug.
- A low dose of meloxicam is maintained to prevent renal stress; ensuring hydration is crucial.
- Guidance for owner handling is included with the prescribed medication (an example is shown on the right).

Stokes Pharmacy
GingiPro™ Duo™
Doxycycline (as monohydrate),
EIDD-1931,
Meloxicam

GingiPro Solo™
EIDD-1931

This is a specially compounded preparation designed for the treatment of cats with Calicivirus-associated stomatitis. It is dispensed using the Topi-CLICK Perl®, a precision dosing applicator that allows accurate, consistent oral doses every time. 1 click = 0.25 mL of paste.

How to Administer

1. Open cap (figure a.): Flip open the top cap on the Topi-CLICK Perl® device.
2. Attach the Snap-on Applicator (figure b.) to the base by aligning and snapping it into the receptor port. Make sure the plunger touches the base.
3. Hold the device vertically and upright.
4. Turn the base until you hear a click, this loads the dose into the applicator (figure c.).
5. Repeat step 4 if another click is required.
6. Remove applicator by gently tilting the applicator off the base – do not pull straight off (figure d.).
7. Wipe the nozzle with a clean tissue and flip the cap back down to close the device.
8. Administer the dose directly into the mouth using the plunger.
9. Gently massage under the chin if needed to encourage swallowing.

Important Notes

- Pregnant women should not handle.
- Wear gloves when handling. Wash hands after administration.
- Store medication at room temperature, away from direct sunlight.
- Wash the applicator thoroughly after use.
- Applicator is dishwasher safe; use with caution to avoid residual medicine from cleaning (max. 30 cycles).
- Follow the dose prescribed by your veterinarian and do not exceed the number of doses.
- This product contains an antibiotic and an antiviral. Use responsibly to help prevent antimicrobial resistance.

Scan QR code: How-To-Use Video

800.754.5222 | stokespharm.com

Scan to view
handout



****Important Notice:** This product is not intended to treat common gingivitis; it is specifically formulated for stomatitis associated with calicivirus. We strongly advise conducting a comprehensive evaluation to determine if the gum inflammation spreads to the palate and the back of the throat before selecting an appropriate prescription or considering alternative treatment options for the situation.

Our Family of Brands



503A Traditional
Compounding Pharmacy



503B FDA Registered
Outsourcing Facility



Stokes/Epicur Online
Ordering Platform